

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau**B16**

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 15/11, C07K 14/00	A2	(11) International Publication Number: WO 96/13583 (43) International Publication Date: 9 May 1996 (09.05.96)
(21) International Application Number: PCT/EP95/04117 (22) International Filing Date: 20 October 1995 (20.10.95) (30) Priority Data: 94116558.1 20 October 1994 (20.10.94) EP (34) Countries for which the regional or international application was filed: DE et al. (71) Applicant (for all designated States except US): MORPHOSYS GESELLSCHAFT FÜR PROTEINOPTIMIERUNG MBH [DE/DE]; Frankfurter Ring 193a, D-80807 Munich (DE). (72) Inventors; and (75) Inventors/Applicants (for US only): PACK, Peter [DE/DE]; Franz-Wolter-Strasse 4/III, D-81925 Oberföhring (DE). LU- PAS, Andrei [DE/DE]; Fasanenstrasse 68a, D-82008 Unter- haching (DE). (74) Agent: VOSSIUS & PARTNER; P.O. Box 86 07 67, D-81634 Munich (DE).		(81) Designated States: JP, US, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>Without international search report and to be republished upon receipt of that report.</i>
(54) Title: TARGETED HETERO-ASSOCIATION OF RECOMBINANT PROTEINS TO MULTI-FUNCTIONAL COMPLEXES (57) Abstract The present invention relates to a method for targeted assembly of distinct active peptide or protein domains into a single complex and to such complexes. The invention relates particularly to the fusion of peptide or protein domains to complementary association domains which are derived from a single tertiary or quaternary structure by segmentation. The association domains are designed to assemble in a complementary fashion, thereby providing multifunctional (poly)peptides.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

TARGETED HETERO-ASSOCIATION OF RECOMBINANT PROTEINS TO MULTI-FUNCTIONAL COMPLEXES

Background of the Invention

Increasingly, there is a need for proteins which combine two or more functions, such as binding or catalysis, in a single structure. Typically, proteins which combine two or more functions are prepared either as fusion proteins or through chemical conjugation of the component functional domains. Both of these approaches suffer from disadvantages. Genetic "single chain" fusions suffer the disadvantages that (i) only a few (2-3) proteins can be fused (Rock et al., 1992, *Prot. Eng.* 5, 583-591), (ii) mutual interference between the component domains may hinder folding, and (iii) the size of the fusion protein may make it difficult to prepare. The alternative, chemical cross-linking *in vitro* following purification of independently expressed proteins, is difficult to control and invariably leads to undefined products and to a severe loss in yield of functional material.

Recently, methods for achieving non-covalent association of two or more of the same functional domains have been developed. This can be achieved through the use of domains attached to peptides which self-associate to form homo-multimers (Pack & Plückthun, 1992, *Biochemistry* 31, 1579-1584). For example, the association of two separately expressed scFv antibody fragments by C-terminally fused amphipathic helices *in vivo* provides homo-dimers of antibody fragments in *E. coli* (PCT/EP93/00082; Pack et al., 1993, *Bio/Technology* 11, 1271-1277) or homo-tetramers; (Pack et al., 1995, *J. Mol. Biol.*, 246, 28-34).

To assemble distinct protein functions such as two antibody fragments with different specificities fused to such association domains, the helices must have a tendency to form hetero-multimers. In principle, this could be achieved with complementary helices such as the hetero-dimerizing JUN and FOS zippers of the AP-1 transcription factor (O'Shea et al., 1992, *Cell* 68, 699-708). The clear disadvantage of association domains based on hetero-associating helices,

however, is their pseudo-symmetry and their similar periodicity of hydrophobic and hydrophilic residues. This structural similarity results in a strong tendency to form homo-dimers and, thus, to lower significantly the yield of hetero-dimers (O'Shea et al., 1992, *Cell* 68, 699-708; Pack, 1994, Ph. D. thesis, Ludwig-Maximilians-Universität München). Furthermore, the formation of JUN/FOS hetero-dimers is kinetically disfavoured and requires a temperature-dependent unfolding of the kinetically favoured homo-dimers, especially JUN/JUN homo-dimers (PCT/EP93/00082; O'Shea et al., 1992, *Cell* 68, 699-708; Pack, 1994, Ph. D. thesis, Ludwig-Maximilians-Universität München). Because of the need for additional purification steps to separate the unwanted homo-dimers from hetero-dimers and the resulting decrease in yield, hetero-association domains based on amphipathic helices do not result in practical advantages compared to conventional chemical coupling.

These disadvantages of the prior art are overcome by the present invention which provides multi-functional polypeptides and methods for the preparation of these multi-functional proteins. This is achieved via the use of association domains which are designed to associate predominantly in a complementary fashion, and not to self-associate.

Detailed Description of the Invention

In the earliest steps of protein folding, peptide chains form a disordered hydrophobic core by collapsing hydrophobic residues into the interior of an intermediate "molten globule". This hydrophobic effect is considered to be the most important driving force of folding (Matthews, 1993, *Annu. Rev. Biochem.* 62, 653 - 683; Fersht, 1993, *FEBS Letters* 325, 5 - 16). The burial of hydrophobic residues and the resulting exclusion of solvent is the determining factor in the stability of compact tertiary structures such as acyl-phosphatase (Pastore et al, *J. Mol. Biol.* 224, 427-440, 1992) interleukin-2 (Brandhuber et al., 1987, *Science* 238, 1707 - 1709), calbindin (Parmentier, 1990, *Adv. Exp. Med. Biol.* 269, 27-34) or ubiquitin (Briggs & Roder, 1992, *Proc. Natl. Acad. Sci. USA* 89, 2017 - 2021).

This concept forms the basis of the present invention, which provides individually encoded peptides or "segments" which, in a single continuous chain, would comprise a compact tertiary structure with a highly hydrophobic core. The component peptides are chosen so as to be asymmetric in their assumed structure, so as not to self-associate to form homo-multimers, but rather to associate in a complementary fashion, adopting a stable complex which resembles the parent tertiary structure. On the genetic level, these segments are encoded by interchangeable cassettes with suitable restriction sites. These standardized cassettes are fused C- or N-terminally to different recombinant proteins via a linker or hinge in a suitable expression vector system.

Thus, the present invention relates to a multi-functional polypeptide comprising:

- (a) a first amino acid sequence attached to at least one functional domain;
- (b) a second amino acid sequence attached to at least one further functional domain; and
- (c) optionally, further amino acid sequences each attached to at least one further functional domain;

wherein any one or more of said amino acid sequences interacts with at least one of said amino acid sequences in a complementary fashion to form a parental, native-like tertiary or optionally quaternary structure and wherein the parental, native-like tertiary or optionally quaternary structure is derived from a single parent polypeptide. In this context, the term parent polypeptide refers to a polypeptide which has a compact tertiary or quaternary structure with a hydrophobic core. The invention provides for many different parent polypeptides to be used as the basis for the association domain. Suitable polypeptides can be identified by searching for compact, single-domain proteins or protein fragments in the database of known protein structures (Protein Data Bank, PDB) and selecting structures that are stable and can be expressed at high yields in recombinant form. These structures can then be analyzed for hydrophobic sub-clusters by the method of Karpeisky and Ilyn (1992, *J. Mol. Biol.* 224, 629-638) or for structural units (such as β -elements or helical hairpin structures) by standard molecular modelling techniques. In a further embodiment, the present invention

provides for multi-functional polypeptides wherein the single parent polypeptide is taken from the list ubiquitin, acyl phosphatase, IL-2, calbindin and myoglobin.

In a preferred embodiment, the present invention provides a multi-functional polypeptide comprising two or more amino acid sequences each attached to at least one functional domain, wherein any two or more of said amino acid sequences can associate in a complementary fashion to provide a parental, native like, tertiary or optionally quaternary structure.

Once structural sub-domains are identified, the protein is dissected in such a way these sub-domains remain intact. The selection process can be expanded to proteins for which no structure is available but which satisfy the criteria of stability and good expression. For these proteins, folding sub-domains can be determined by hydrogen exchange pulse-labelling of backbone amides during the folding reaction, followed by NMR detection in the native state (Roder et al., 1988, *Nature* 355, 700-704; Udgaonkar & Baldwin, 1988, *Science* 255, 594-597). Alternatively, folding sub-domains can be identified by mild proteolysis, denaturation, purification of fragments and reconstitution *in vitro* (Tasayco & Carey, 1992, *Science* 255, 594-597; Wu et al., 1993, *Biochemistry* 32, 10271-10276). Finally, additional clues for the choice of cleavage sites can be obtained from the exon structure in the case of eukaryotic proteins, since the exons frequently (though not always) correspond to structural sub-domains of a protein. This has, for example, been discussed for the case of myoglobin (Go 1981, *Nature* 291, 90).

The yield of properly assembled molecules is expected to decrease significantly for constructs in which a protein domain is divided into three or more parts. This is due to the fact that several sub-domains must come together simultaneously to form a viable structure. This effect is countered by dividing the polypeptide chain into sub-domains that represent folding units (identified by the methods described above). Thus, not only the final, assembled complex but also assembly intermediates will have the stability necessary to allow their accumulation in the

host during expression, resulting in a greatly improved kinetic behaviour of the system.

In solution, the isolated segments have little secondary structure and remain monomeric or form transient, non-specific and easily disrupted aggregates. Only upon mixing, either by separate expression and purification, or by co-expression, can the concerted folding of complementary segments provide the necessary intermediate interaction of residues (Matthews, 1993, *Annu. Rev. Biochem.* 62, 653 - 683) that results in the formation of a compact, native-like structure. This association, mainly driven by the burial of hydrophobic residues of all segments into a single hydrophobic core, leads to a targeted assembly of the N- or C-terminally fused proteins to a multi-functional complex *in vivo* or *in vitro*.

Optionally, the reconstituted native-like structure may also contribute an enzymatic or binding activity to increase the number of effector functions in the assembled complex. Accordingly, the present invention also provides a multi-functional polypeptide as described above, in which the native-like, tertiary or quaternary structure provides a biological activity. For example, when acyl phosphatase is used as the basis of the association domain, it is expected that the multi-functional polypeptide will retain some phosphatase activity.

The present invention provides for many different types of functional domains to be linked into the multi-functional polypeptide. Particularly preferred are cases in which one or more, preferably two, of said functional domains are fragments derived from molecules of the immunoglobulin superfamily. In particularly preferred embodiments, said fragments are antibody fragments. Also preferred are cases in which at least one of the functional domains possesses biological activity other than that associated with a fragment derived from a member of the immunoglobulin superfamily. By way of example, the present invention provides for the targeted assembly of enzymes, toxins, cytokines, peptide hormones, immunoglobulins, metal binding domains, soluble receptors, lectins, lipoproteins, purification tails and bioactive peptides to multi-functional complexes (Fig. 1) based on a modular system of expression vectors, restriction sites and "plug-in" gene cassettes coding for assembly segments, peptide linkers and functional domains (Fig.2).

If covalent linkage between the segments is necessary to prevent dissociation at low concentrations, cysteines can be introduced to form inter-segmental disulphide bridges between the amino acid sequences which comprise the association domain (Ecker et al., 1989, *J. Biol. Chem.* 264, 1887-1893; Pack & Plückthun, 1992, *Biochemistry* 31, 1579-1584). Accordingly, the present invention provides multi-functional polypeptides wherein the folding of the component amino acid sequences is stabilized by a covalent bond.

In order to provide some flexibility between the association domain and the appended functional domains, it may be desired to incorporate a linker peptide. Accordingly, the present invention provides for multi-functional polypeptides of the type described above wherein at least one of the functional domains is coupled to said amino acid sequence via a flexible peptide linker. By way of example, the flexible linker may be derived from the hinge region of an antibody. The invention enables even more complex multi-functional polypeptides to be constructed via the attachment of at least one further (poly)peptide to one or more of said amino acid sequences. By way of example, the further (poly)peptide can be taken from the list enzymes, toxins, cytokines, peptide hormones, immunoglobulins, metal binding domains, soluble receptors, lectins, lipoproteins, purification tails, in particular peptides which are able to bind to an independent binding entity, bioactive peptides, preferably of 5 to 15 amino acid residues, metal binding proteins, DNA binding domains, transcription factors and growth factors.

For therapeutic purposes, it is often desirable that proteinaceous substances display the minimum possible immunogenicity. Accordingly, the present invention provides for multi-functional polypeptides as described above in which at least one of said amino acid sequences, functional domains, or further (poly)peptides is of human origin.

In addition to the peptides and proteins provided above, the present invention also provides for DNA sequences, vectors, preferably bicistronic vectors, vector cassettes, characterised in that they comprise a DNA sequence encoding an amino acid sequence and optionally at least one further (poly)peptide comprised in the multifunctional polypeptide of the invention, and additionally at least one,

preferably singular cloning sites for inserting the DNA encoding at least one further functional domain or that they comprise DNA sequences encoding the amino acid sequences, and optionally the further (poly)peptide(s) comprised in the multifunctional polypeptide of the invention and suitable restriction sites for the cloning of DNA sequences encoding the functional domains, such that upon expression of the DNA sequences after the insertion of the DNA sequences encoding the functional domains into said restriction sites, in a suitable host the multifunctional polypeptide of the invention is formed. In a preferred embodiment said vector cassette is characterised in that it comprises the inserted DNA sequence(s) encoding said functional domain(s) and host cells transformed with at least one vector or vector cassette of the invention which can be used for the preparation of said multi-functional polypeptides.

In a further preferred embodiment, said host cell is a mammalian, preferably human, yeast, insect, plant or bacterial, preferably *E. coli* cell.

The invention further provides for a method for the production of a multifunctional polypeptide of the invention, which comprises culturing the host cell of the invention in a suitable medium, and recovering said multifunctional polypeptide produced by said host cell.

In a further embodiment, the invention relates to a method for the production of a multifunctional polypeptide of the invention which comprises culturing at least two host cells of the invention in a suitable medium, said host cells each producing only one of said first and said second amino acid sequences attached to at least one further functional domain, recovering the amino acid sequences, mixing thereof under mildly denaturing conditions and allowing in vitro folding of the multifunctional polypeptide of the invention from said amino acid sequences.

In a particular preferred embodiment, said method is characterised in that the further amino acid sequences attached to at least one further functional domain

are/is produced by at least one further host cell not producing said first or second amino acid sequence.

In another particularly preferred embodiment of the invention, said method is characterised in that at least one further amino acid sequence attached to at least one further functional domain is produced by the host cell of the invention producing said first or second amino acid sequence.

In further preferred embodiments, the present invention provides for pharmaceutical and diagnostic compositions comprising the multi-functional polypeptides described above, said pharmaceutical compositions optionally comprising a pharmaceutically acceptable carrier. Finally, the invention provides for a kit comprising one or more vector cassettes useful in the preparation of said multi-functional polypeptides.

The invention is now illustrated by reference to the following examples, which are provided for the purposes of illustration only and are not intended to limit the scope of the invention.

Example 1: Segmented human ubiquitin as an assembly device

Ubiquitin is a compact intracellular protein of only 76 residues (Fig. 3) and a molecular weight of 5 kDa. It shows the highest conservation among all known proteins and is involved in the degradation pathway of intracellular eukaryotic proteins by forming intermediate isopeptide bonds to its C-terminus and to Lys48 (Hershko & Ciechanover, 1992, *Ann. Rev. Biochem.* 61, 761-807).

To use ubiquitin as an assembly device, the unwanted function can be abolished by truncation of the last three C-terminal residues (--Arg-Gly-Gly), and the exchange of Lys48 to Arg, which prevents the formation of isopeptide bonds to this residue. The altered sequence is then divided in a loop at position Gly36, so that the hydrophobic core falls apart into two segments (called ALPHA and BETA). The synthetic nucleotide sequence of the segments (Fig 4, 5) carry appropriate restriction sites (MroI-HindIII) at the termini, so that the cassette encoding the segments can be easily ligated to a EcoRI-MroI cassette encoding the flexible linker (hinge of hulgG3; Fig. 6). The cassettes are inserted into the expression vector pIG3 (EcoRI-HindIII; Fig. 7) encoding the scFv fragment of the antibody McPC603 under the lac promoter/operator (Ge et al., 1995, in: *Antibody engineering: A practical approach*. IRL Press, New York, Borrebaeck ed., 229-261). Insertion of a second functional fragment (scFv fragment of the anti- β -lactam antibody 2H10 with phoA signal sequence) linked to association segment BETA as an XbaI-HindIII DNA fragment (Fig. 8) results in a di-cistronic expression vector (Pack, 1994, Ph. D. thesis, Ludwig-Maximilians-Universität München). After induction with IPTG and translation, the signal sequences guide the antibody fragments fused to the assembly segments to the periplasm, where they assemble to a complex with a reconstituted native-like ubiquitin fold and two different antibody specificities. The complex, a bispecific immunoglobulin, can be recovered and purified by affinity chromatography of cell extract (Pack, 1994, Ph. D. thesis, Ludwig-Maximilians-Universität München).

Example 2: Covalent linkage of the native-like tertiary structure of the assembly device by engineered disulphide bridges and combination of a C-terminal peptide linker with an in-frame restriction site.

The conformational stability of undivided, native ubiquitin can be enhanced by introduction of disulfides at positions 4 and 66 without perturbation in the backbone (Ecker et al., 1989, *J. Biol. Chem.* 264, 1887-1893; Fig. 9). In the context of this invention, the engineering of disulfide bridges provides the covalent linkage of segments (Fig. 10, 11) after co-folding and assembly.

To raise the number of possible functional domains in the assembled complex, a C-terminal peptide can be fused to one or more of the segments of the assembly device. To fuse a functional domain like an enzyme, cytokine, antibody fragment, purification peptide or toxin to this linker, a restriction site, preferably unique, has to be introduced in-frame (Fig. 11). Gene synthesis, cloning, expression as well as recovery of the assembled, covalently linked complex is according to example 1.

Example 3: Segmented human interleukin-2 (IL2) as an assembly device

Human interleukin-2 (Brandhuber et al., 1987, *Science* 238, 1707 - 1709; Kuziel & Greene, 1991, in: *The Cytokine Handbook*. Academic Press, 84-100) is used as an assembly device by segmentation between position His79 and Lys 80 (Fig. 12). The device, encoded by MroI-Ascl-HindII gene cassettes (Fig. 13, 14) combines the low immunogenicity of the plasmatic protein with a preferable effector function of the native-like cytokine structure and an inter-segmental cysteine bridge (Cys58-Cys105) after assembly. The combination of one or more antibody fragments against tumor antigens with additional cytokines like IL6 or

IL12 targets the multi-cytokine complex (Rock et al., 1992, *Prot. Eng.* 5, 583-591) directly to the tumour.

Example 4 : Segmented human apomyoglobin as an assembly device with three segments

To use more than two segments of a native structure as an assembly device, the hydrophobic interface between the segments has to be large enough to provide the sufficient hydrophobic interaction for non-covalent linkage. Myoglobin (Fig.15) is expressible in large amounts in *E. coli* (Guillemette et al., 1991, *Protein Eng.* 4, 585-592). Up to six functional domains can be assembled by a threefold segmented structure (Fig. 16, 17, 18), three at the N-termini and three at the C-termini of the segments. The presence of heme additionally stabilizes the native-like apomyoglobin fold and can be used as a switch to influence the association constant of the multi-functional complex.

Example 5: Bioactive peptides as functional domains

Certain peptides derived from amphipathic loop structures of LPS-binding proteins (Hoess et al., 1993, *EMBO J.* 12, 3351-3356) are able to neutralize endotoxin. This effect is enhanced by multivalent display of these short peptides (10-15 residues; Hoess, unpublished results). The present invention provides a method to express and assemble several of short peptides (Fig.19), fused to an assembly segment, in a multivalent complex or in combination with other functional domains. The peptides can be fused either to the N-or to the C-terminus (Fig. 20, 21) of the assembly domain via the peptide linkers.

Example 6: A purification tail for IMAC as a functional domain

Peptide tails consisting of histidines are able to coordinate metal ions. They are used for purification of native proteins in immobilized metal affinity chromatography (IMAC). Multivalent display of the purification tail considerably improves the maximum purity achievable by IMAC (Lindner et al., 1992, *Methods: a companion to methods in enzymology* 4, 41-56). One or more gene cassettes (Fig. 22) encoding a polyhistidine tail can be fused to the assembly segment to provide a simple and efficient purification method for multi-functional complexes.

Example 7: The platelet aggregation inhibitor decorsin as a functional domain

Decorsin, a 39 residue protein of the leech *Macrobdella decora* (Fig. 23), acts as a potent antagonist of the platelet glycoprotein IIb-IIIa (Seymour et al., 1990, *J. Biol. Chem.* 265, 10143-10147). The gene cassette encoding the decorsin can be fused C- or N-terminally to an association segment (Fig. 24, 25). In arterial thrombotic diseases, a multivalent decorsin complex combined with an anti-fibrin antibody fragment can act as a powerful antithrombotic agent.

Claims

1. A multifunctional polypeptide comprising:
 - (a) a first amino acid sequence attached to at least one functional domain;
 - (b) a second amino acid sequence attached to at least one further functional domain; and
 - (c) optionally, further amino acid sequences each attached to at least one further functional domain;wherein any one or more of said amino acid sequences interacts with at least one of said amino acid sequences in a complementary fashion to form a parental, native-like tertiary or optionally quaternary structure and wherein said parental, native-like tertiary or optionally quaternary structure is derived from a single parent polypeptide.
2. The multifunctional polypeptide according to claim 1, wherein said single parent polypeptide is ubiquitin, acyl-phosphatase, IL2, calbindin or apomyoglobin.
3. The multifunctional polypeptide according to claim 1 or 2, wherein said parental, native-like tertiary or optionally quaternary structure is biologically active.
4. The multifunctional polypeptide according to any one of claims 1 to 3, wherein at least one of said functional domains is a fragment derived from a member of the immunoglobulin superfamily.
5. The multifunctional polypeptide according to claim 4, wherein two of said functional domains are fragments derived from members of the immunoglobulin superfamily.


6. The multifunctional polypeptide according to claim 4 or 5, wherein said fragments are antibody fragments.
7. The multifunctional polypeptide according to any one of claims 1 to 6, wherein at least one of said functional domains is a biologically active molecule or a derivative thereof other than a fragment derived from a member of the immunoglobulin superfamily.
8. The multifunctional polypeptide according to any one of claims 1 to 6, wherein the folding of the amino acid sequences is stabilised by a covalent bonding.
9. The multifunctional polypeptide according to any one of claims 1 to 8, wherein at least one of said functional domains is coupled to said amino acid sequence(s) via a flexible peptide linker.
10. The multifunctional polypeptide according to claim 9, wherein said flexible peptide linker is an antibody hinge region.
11. The multifunctional polypeptide according to any one of claims 1 to 10, wherein at least one of said amino acid sequences is coupled to at least one further (poly)peptide.
12. The multifunctional polypeptide according to claim 11, wherein said further (poly)peptide is an enzyme, a toxin, a cytokine, a metal binding site, a metal binding protein, a soluble receptor, a DNA-binding domain, a transcription factor, an immunoglobulin, a bioactive peptide of 5 to 15 amino acid residues, a peptide hormone, a growth factor, a lectin, a lipoprotein, and a peptide which is able to bind to an independent binding entity.

13. The multifunctional polypeptide according to any one of claims 1 to 12, wherein at least one of said amino acid sequences, functional domains or further (poly)peptide(s) is of human origin.
14. A DNA sequence encoding an amino acid sequence and at least one functional domain and, optionally, at least one further functional (poly)peptide comprised in the multifunctional polypeptide of any one of claims 1 to 13.
15. A vector comprising at least one DNA molecule of claim 14.
16. The vector of claim 15, which is a bicistronic vector.
17. A vector cassette characterised in that it comprises a DNA sequence encoding an amino acid sequence and optionally at least one further (poly)peptide comprised in the multifunctional polypeptide of any one of claims 1 to 13, and additionally at least one, preferably a singular cloning site for inserting the DNA encoding at least one further functional domain.
18. A vector cassette characterised in that it comprises DNA sequences encoding the amino acid sequences, and optionally the further (poly)peptide(s) comprised in the multifunctional polypeptide of any one of claims 1 to 13, and suitable restriction sites for the cloning of DNA sequences encoding the functional domains, such that upon expression of the DNA sequences after the insertion of the DNA sequences encoding the functional domains into said restriction sites, in a suitable host the multifunctional polypeptide according to any one of claims 1 to 13 is formed.

19. The vector cassette according to claim 17 or 18 characterised in that it comprises the inserted DNA sequence(s) encoding said functional domain(s).
20. A host cell transformed with at least one vector according to claim 15 or 16, or at least one vector cassette according to claim 19.
21. The host cell according to claim 20, which is a mammalian, preferably human, yeast, insect, plant or bacterial, preferably E. coli cell.
22. A method for the production of a multifunctional polypeptide according to any one of claims 1 to 13, which comprises culturing the host cell according to claim 20 or 21 in a suitable medium, and recovering said multifunctional polypeptide produced by said host cell.
23. A method for the production of a multifunctional polypeptide according to any one of claims 1 to 13 which comprises culturing at least two host cells according to claim 20 or 21 in a suitable medium, said host cells each producing only one of said first and said second amino acid sequences attached to at least one further functional domain, recovering the amino acid sequences, mixing thereof under mildly denaturing conditions and allowing in vitro folding of the multifunctional polypeptide according to any one of claims 1 to 13 from said amino acid sequences.
24. The method according to claim 23, wherein the further amino acid sequence(s) (each) attached to at least one further functional domain are/is produced by at least one further host cell not producing said first or second amino acid sequence.

25. The method according to claim 23, wherein at least one further amino acid sequence attached to at least one further functional domain is produced by the host cell according to claim 20 or 21 producing said first or second amino acid sequence.
26. A pharmaceutical composition comprising the multifunctional polypeptide according to any one of claims 1 to 13 optionally in combination with a pharmaceutically acceptable carrier.
27. A diagnostic composition comprising the multifunctional polypeptide according to any one of claims 1 to 13.
28. A kit comprising at least one vector cassette according to claim 17 or 18.


1/15

Fig.1: segmented tertiary structure for a targeted hetero-association

RECTIFIED SHEET (RULE 91)

ISA/EP

Fig. 2 : Modular cistron encoding functional domains
N- and/or C-terminally fused to the assembly device



3/15

Fig. 3 : protein sequence of human ubiquitin (segmented after Gly35)

```

1      10      20      30      40      50
MQIFVKLTG KTITLEVEPS DTIENVKAKI QDKEGIPPDQ QRLIFAGKQL

      60      70
EDGRTLSDYN IQKESTLHLV LRLRGG**

```

Fig. 4: MroI-Hind III gene cassette encoding for segment ALPHA of ubiquitin

MroI

```

S  G  M  Q  I  F  V  K  T  L  T  G  K  T  I  T  L  E
TCC GGA ATG CAG ATC TTC GTT AAA ACC CTG ACC GGT AAA ACC ATC ACC CTG GAA
      9      18      27      36      45      54
AGG CCT TAC GTC TAG AAG CAA TTT TGG GAC TGG CCA TTT TGG TAG TGG GAC CTT

```

```

V  E  P  S  D  T  I  E  N  V  K  A  K  I  Q  D  K  E
GTT GAA CCG TCT GAC ACC ATC GAA AAC GTT AAA GCT AAA ATC CAG GAC AAA GAA
      63      72      81      90      99      108
CAA CTT GGC AGA CTG TGG TAG CTT TTG CAA TTT CGA TTT TAG GTC CTG TTT CTT

```

HindIII

```

G  *  *  A
GGT TGA TAA GCT T 3'
      117
CCA ACT ATT CGA A 5'

```

4/15

Fig. 5: MroI-Hind III gene cassette encoding for segment BETA of ubiquitin

MroI

S	G	I	P	P	D	Q	Q	R	L	I	F	A	G	R	Q	L	E
TCC	GGA	ATC	CCG	CCG	GAC	CAG	CAG	CGT	CTG	ATC	TTC	GCT	GGT	CGT	CAG	CTG	GAA
		9			18			27			36			45			54
AGG	CCT	TAG	GGC	GGC	CTG	GTC	GTC	GCA	GAC	TAG	AAG	CGA	CCA	GCA	GTC	GAC	CTT

D	G	R	T	L	S	D	Y	N	I	Q	K	E	S	T	L	H	L
GAC	GGT	CGT	ACC	CTG	TCT	GAC	TAC	AAC	ATC	CAG	AAA	GAA	TCT	ACC	CTG	CAC	CTG
		63			72			81			90			99			108
CTG	CCA	GCA	TGG	GAC	AGA	CTG	ATG	TTG	TAG	GTC	TTT	CTT	AGA	TGG	GAC	GTG	GAC

HindIII


V	L	R	L	*	*	
GTT	CTG	CGT	CTG	TGA	TAA	3'
		117			126	
CAA	GAC	GCA	GAC	ACT	ATT	5'

Fig. 6: EcoRI-MroI gene cassette encoding a flexible linker (huIgG3)

EcoRI										MroI				
	E	F	T	P	L	G	D	T	T	H	T	S	G	
5'	GAA	TTC	ACC	CCG	CTG	GGT	GAC	ACC	ACC	CAC	ACC	TCC	GGA	3'
			9			18			27			36		
3'	CTT	AAG	TGG	GGC	GAC	CCA	CTG	TGG	TGG	GTG	TGG	AGG	CCT	5'

5/15

Fig. 7 : Construction of monocistronic expression vector




RECTIFIED SHEET (RULE 91)

ISA/EP

6/15

Fig. 8: Construction of dicistronic co-expression vectors



RECTIFIED SHEET (RULE 91)

ISA/EP

7/15

Fig. 9 : protein sequence of human ubiquitin with intersegmental disulfides Cys4 and Cys66 (segmented after Gly35)

```

1      10      20      30      40      50
MQICVKLTG KTITLEVEPS DTIENVKAKI QDKEGIPPDQ QRLIFAGKQL

      60      70
EDGRTLSDYN IQKESCLHLV LRLRGG**

```

Fig. 10: MroI-Hind III gene cassette encoding for segment ALPHA-CYS4 of ubiquitin

MroI

```

S  G  M  Q  I  C  V  K  T  L  T  G  K  T  I  T  L  E
TCC GGA ATG CAG ATC TGC GTT AAA ACC CTG ACC GGT AAA ACC ATC ACC CTG GAA
      9      18      27      36      45      54
AGG CCT TAC GTC TAG ACG CAA TTT TGG GAC TGG CCA TTT TGG TAG TGG GAC CTT

V  E  P  S  D  T  I  E  N  V  K  A  K  I  Q  D  K  E
GTT GAA CCG TCT GAC ACC ATC GAA AAC GTT AAA GCT AAA ATC CAG GAC AAA GAA
      63      72      81      90      99      108
CAA CTT GGC AGA CTG TGG TAG CTT TTG CAA TTT CGA TTT TAG GTC CTG TTT CTT

```

HindIII

```

G  *  *  A
GGT TGA TAA GCT T 3'
      117
CCA ACT ATT CGA A 5'

```

8/15

Fig. 11 : MroI-AscI-Hind III gene cassette encoding for segment BETA-CYS66 with C-terminal GSGGAP linker of ubiquitin

MroI

S G I P P D Q Q R L I F A G R Q L E
 TCC GGA ATC CCG CCG GAC CAG CAG CGT CTG ATC TTC GCT GGT CGT CAG CTG GAA
 9 18 27 36 45 54
 AGG CCT TAG GGC GGC CTG GTC GTC GCA GAC TAG AAG CGA CCA GCA GTC GAC CTT

D G R T L S D Y N I Q K E S C L H L
 GAC GGT CGT ACC CTG TCT GAC TAC AAC ATC CAG AAA GAA TCT TGC CTG CAC CTG
 63 72 81 90 99 108
 CTG CCA GCA TGG GAC AGA CTG ATG TTG TAG GTC TTT CTT AGA ACG GAC GTG GAC

 AscI HindIII
 V L R L G G S G G A P * *
 GTT CTG CGT CTG GGG GGG AGC GGA GGC GCG CCG TGA TAA 3'
 117 126
 CAA GAC GCA GAC CCC CCC TCG CCT CCG CGC GGC ACT ATT 5'

Fig. 12: Protein sequence of human IL-2 (segmented after His79)

10 20 30 40 50 60
 APTSSSTKKT QLQLEHLLLD LQMILNGINN YKNPKLTRML TFKFYMPKKA TELKHLQCLE

 70 80 90 100 110 120
 EELKPLEEVL NLAQSKNFHL RPRDLISNIN VIVLELKGSE TTFMCEYADE TATIVEFLNR

 130
 WITFCQSIIS TLT

MroI																			
S	G	A	P	T	S	S	S	T	K	K	T	Q	L	Q	L	E	H		
TCC	GGA	GCA	CCT	ACT	TCA	AGT	TCT	ACA	AAG	AAA	ACA	CAG	CTA	CAA	CTG	GAG	CAT		
		9			18			27			36			45			54		
AGG	CCT	CGT	GGA	TGA	AGT	TCA	AGA	TGT	TTC	TTT	TGT	GTC	GAT	GTT	GAC	CTC	GTA		
L	L	L	D	L	Q	M	I	L	N	G	I	N	N	Y	K	N	P		
TTA	CTG	CTG	GAT	TTA	CAG	ATG	ATT	TTG	AAT	GGA	ATT	AAT	AAT	TAC	AAG	AAT	CCC		
		63			72			81			90			99			108		
AAT	GAC	GAC	CTA	AAT	GTC	TAC	TAA	AAC	TTA	CCT	TAA	TTA	TTA	ATG	TTC	TTA	GGG		
K	L	T	R	M	L	T	F	K	F	Y	M	P	K	K	A	T	E		
AAA	CTC	ACC	AGG	ATG	CTC	ACA	TTT	AAG	TTT	TAC	ATG	CCC	AAG	AAG	GCC	ACA	GAA		
		117			126			135			144			153			162		
TTT	GAG	TGG	TCC	TAC	GAG	TGT	AAA	TTC	AAA	ATG	TAC	GGG	TTC	TTC	CGG	TGT	CTT		
L	K	H	L	Q	C	L	E	E	E	L	K	P	L	E	E	V	L		
CTG	AAA	CAT	CTT	CAG	TGT	CTA	GAA	GAA	GAA	CTC	AAA	CCT	CTG	GAG	GAA	GTG	CTA		
		171			180			189			198			207			216		
GAC	TTT	GTA	GAA	GTC	ACA	GAT	CTT	CTT	CTT	GAG	TTT	GGA	GAC	CTC	CTT	CAC	GAT		
														AscI		HindIII			
N	L	A	Q	S	K	N	F	H	G	G	S	G	G	A	P	*			
AAT	TTA	GCT	CAA	AGC	AAA	AAC	TTT	CAC	GGG	GGG	AGC	GGA	GGC	CGC	CCG	TGA	T		
		225			234			243			252			261					
TTA	AAT	CGA	GTT	TCG	TTT	TTG	AAA	GTG	CCC	CCC	TCG	CCT	CCG	CGC	GGC	ACT	A		

10/15

Fig. 14 : MroI-AscI-Hind III gene cassette encoding for segment BETA of human IL-2

MroI
 S G L R P R D L I S N I N V I V L E
 TCC GGA TTA AGA CCC AGG GAC TTA ATC AGC AAT ATC AAC GTA ATA GTT CTG GAA
 9 18 27 36 45 54
 AGG CCT AAT TCT GGG TCC CTG AAT TAG TCG TTA TAG TTG CAT TAT CAA GAC CTT

L K G S E T T F M C E Y A D E T A T
 CTA AAG GGA TCT GAA ACA ACA TTC ATG TGT GAA TAT GCT GAT GAG ACA GCA ACC
 63 72 81 90 99 108
 GAT TTC CCT AGA CTT TGT TGT AAG TAC ACA CTT ATA CGA CTA CTC TGT CGT TGG

I V E F L N R W I T F C Q S I I S T
 ATT GTA GAA TTT CTG AAC AGA TGG ATT ACC TTT TGT CAA AGC ATC ATC TCA ACA
 117 126 135 144 153 162
 TAA CAT CTT AAA GAC TTG TCT ACC TAA TGG AAA ACA CTT TCG TAG TAG AGT TGT

 AscI HindIII
 L T G G S G G A P *
 CTG ACT GGG GGG AGC GGA GGC GCG CCG TGA T 3'
 171 180 189
 GAC TGA CCC CCC TCG CCT CCG CGC GGC ACT A 5'

11/15

Fig. 15 Protein sequence of human apomyoglobin (cut after Lys47 and Lys98)

```

                                ✂
mglsdgewql vlnvwgkvea dipghgqevl irlfkghpet lekfdkfkhl
51
ksedemkase dlkkhgvatl talggilkkk ghheaeikpl aqshatkxki
101
pvkylefise ciqvqlsqkh pgdfgadaeg amnkalelfr kdmasykel
151
gfqg

```

Fig. 16 : MroI-AscI-Hind III gene cassette encoding for segment ALPHA of human apomyoglobin

MroI

S	G	M	G	L	S	D	G	E	W	Q	L	V	L	N	V	W	G
TCC	GGA	ATG	GGT	CTG	TCT	GAC	GGT	GAA	TGG	CAG	CTG	GTT	CTG	AAC	GTT	TGG	GGT
		9			18			27			36			45			54
AGG	CCT	TAC	CCA	GAC	AGA	CTG	CCA	CTT	ACC	GTC	GAC	CAA	GAC	TTG	CAA	ACC	CCA

K	V	E	A	D	I	P	G	H	G	Q	E	V	L	I	R	L	F
AAA	GTT	GAA	GCT	GAC	ATC	CCG	GGT	CAC	GGT	CAG	GAA	GTT	CTG	ATC	CGT	CTG	TTC
		63			72			81			90			99			108
TTT	CAA	CTT	CGA	CTG	TAG	GGC	CCA	GTG	CCA	GTC	CTT	CAA	GAC	TAG	GCA	GAC	AAG

K	G	H	P	E	T	L	E	K	F	D	K	F	K	G	G	S	G
AAA	GGT	CAC	CCG	GAA	ACC	CTG	GAA	AAA	TTC	GAC	AAA	TTC	AAA	GGG	GGG	AGC	GGA
		117			126			135			144			153			162
TTT	CCA	GTG	GGC	CTT	TGG	GAC	CTT	TTT	AAG	CTG	TTT	AAG	TTT	CCC	CCC	TCG	CCT

AscI		HindIII
G	A	P
GGC	CGC	CCG TGA T 3'
		171
CCG	CGC	GGC ACT A 5'

12/15

Fig. 17: MroI-AscI-Hind III gene cassette encoding for segment BETA of human apomyoglobin

MroI
 S G H L K S E D E M K A S E D L K K
 TCC GGA CAC CTG AAA TCT GAA GAC GAA ATG AAA GCA TCT GAA GAC CTG AAA AAA
 9 18 27 36 45 54
 AGG CCT GTG GAC TTT AGA CTT CTG CTT TAC TTT CGT AGA CTT CTG GAC TTT TTT

H G A T V L T A L G G I L K K K G H
 CAC GGT GCT ACC GTT CTG ACC GCT CTG GGT GGT ATC CTG AAA AAA AAA GGT CAC
 63 72 81 90 99 108
 GTG CCA CGA TGG CAA GAC TGG CGA GAC CCA CCA TAG GAC TTT TTT TTT CCA GTG

H E A E I K P L A Q S H A T K H K G
 CAC GAA GCT GAA ATC AAA CCG CTG GCT CAG TCT CAC GCT ACC AAA CAC AAA GGG
 117 126 135 144 153 162
 GTG CTT CGA CTT TAG TTT GGC GAC CGA GTC AGA GTG CGA TGG TTT GTG TTT CCC

 AscI HindIII
 G S G G A P *
 GGG AGC GGA GGC GCG CCG TGA T 3'
 171 180
 CCC TCG CCT CCG CGC GGC ACT A 5'

13/15

Fig.18: MroI-AscI-Hind III gene cassette encoding for segment GAMMA of human apomyoglobin

MroI

S	G	I	P	V	K	Y	L	E	F	I	S	E	C	I	I	Q	V
TCC	GGA	ATC	CCG	GTT	AAA	TAC	CTG	GAG	TTC	ATC	TCT	GAA	TGC	ATC	ATC	CAG	GTT
		9			18			27			36			45			54
AGG	CCT	TAG	GGC	CAA	TTT	ATG	GAC	CTC	AAG	TAG	AGA	CTT	ACG	TAG	TAG	GTC	CAA

L	Q	S	K	H	P	G	D	F	G	A	D	A	E	G	A	M	N
CTG	CAG	TCT	AAA	CAC	CCG	GGT	GAC	TTC	GGT	GCT	GAC	GCT	GAA	GGT	GCT	ATG	AAC
		63			72			81			90			99			108
GAC	GTC	AGA	TTT	GTG	GGC	CCA	CTG	AAG	CCA	CGA	CTG	CGA	CTT	CCA	CGA	TAC	TTG

K	A	L	E	L	F	R	K	D	M	A	S	N	Y	K	E	L	G
AAA	GCT	CTG	GAA	CTG	TTC	CGT	AAA	GAC	ATG	GCT	TCT	AAC	TAC	AAA	GAA	CTG	GGT
		117			126			135			144			153			162
TTT	CGA	GAC	CTT	GAC	AAG	GCA	TTT	CTG	TAC	CGA	AGA	TTG	ATG	TTT	CTT	GAC	CCA
F	S	Q	F	Q	E	T	F	V	H	S	R	V	V	F	F	Q	T

										AscI		HindIII	
F	Q	G	G	G	S	G	G	A	P	*			
TTC	CAG	GGT	GGG	GGG	AGC	GGA	GGC	GCG	CCG	TGA	T	3'	
		171			180			189					
AAG	GTC	CCA	CCC	CCC	TCG	CCT	CCG	CGC	GGC	ACT	A	5'	

14/15

Fig. 19: Peptide sequence of an endotoxin-neutralizing peptide as a functional domain

1 11
RWKVRKSFFKLQ

Fig.20: N-terminal EcoRV-EcoRI cassette encoding an endotoxin-neutralizing peptide

EcoRV														EcoRI			
I	M	R	W	K	V	R	K	S	F	F	K	L	Q	E	F		
5'	ATC	ATG	CGT	TGG	AAA	GTT	CGT	AAA	TCT	TTC	TTC	AAA	CTG	CAG	GAA	TTC	3'
			9			18			27			36			45		
3'	TAG	TAC	GCA	ACC	TTT	CAA	GCA	TTT	AGA	AAG	AAG	TTT	GAC	GTC	CTT	AAG	5'

Fig.21: C-terminal AscI-HindIII cassette encoding an endotoxin-neutralizing peptide

AscI														HindIII			
A	P	R	W	K	V	R	K	S	F	F	K	L	Q	*	*		
5'	GCG	CCG	CGT	TGG	AAA	GTT	CGT	AAA	TCT	TTC	TTC	AAA	CTG	CAG	TGA	TAA	3'
			9			18			27			36			45		
3'	CGC	GGC	GCA	ACC	TTT	CAA	GCA	TTT	AGA	AAG	AAG	TTT	GAC	GTC	ACT	ATT	5'

Fig. 22 AscI-HINDIII Gene cassette encoding a purification tail for IMAC

AscI								Hind III		
A	P	H	H	H	H	H	H	*	*	
5'	GCG	CCG	CAC	CAC	CAC	CAC	CAC	TGA	TAA	3'
			9			18		27		
3'	CGC	GGC	GTG	GTG	GTG	GTG	CAC	ACT	ATT	5'

15/15

Fig. 23 Protein sequence of the platelet aggregation inhibitor decorsin as a functional domain

```

1          11          21          31
APRLPQCQGD DQEKCLCNKD ECPFGQCRFP RGDADPYCE

```

Fig. 24 N-terminal EcoRV-EcoRI cassette encoding the platelet aggregation inhibitor decorsin

```

EcoRV
D I A P R L P Q C Q G D D Q E K C L
GAT ATC GCT CCG CGT CTG CCG CAG TGC CAG GGT GAC GAC CAG GAA AAA TGC CTG
      9      18      27      36      45      54
CTA TAG CGA GGC GCA GAC GGC GTC ACG GTC CCA CTG CTG GTC CTT TTT ACG GAC

C N K D E C P P G Q C R F P R G D A
TGC AAC AAA GAC GAA TGC CCG CCG GGT CAG TGC CGT TTC CCG CGT GGT GAC GCT
      63      72      81      90      99      108
ACG TTG TTT CTG CTT ACG GGC GGC CCA GTC ACG GCA AAG GGC GCA CCA CTG CGA

EcoRI
D P Y C E F
GAC CCG TAC TGC GAA TTC 3'
      117      126
CTG GGC ATG ACG CTT AAG 5'

```

Fig. 25 C-terminal AscI-HindIII cassette encoding the platelet aggregation inhibitor decorsin

```

AscI
A P A P R L P Q C Q G D D Q E K C L
GCG CCG GCT CCG CGT CTG CCG CAG TGC CAG GGT GAC GAC CAG GAA AAA TGC CTG
      12      21      30      39      48      57
CGC GGC CGA GGC GCA GAC GGC GTC ACG GTC CCA CTG CTG GTC CTT TTT ACG GAC

C N K D E C P P G Q C R F P R G D A
TGC AAC AAA GAC GAA TGC CCG CCG GGT CAG TGC CGT TTC CCG CGT GGT GAC GCT
      66      75      84      93      102      111
ACG TTG TTT CTG CTT ACG GGC GGC CCA GTC ACG GCA AAG GGC GCA CCA CTG CGA

HindIII
D P Y C E * *
GAC CCG TAC TGC GAA TGA TAA 3'
      120      129
CTG GGC ATG ACG CTT ACT ATT 5'

```

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶: C12N 15/11, C07K 14/00	A3	(11) International Publication Number: WO 96/13583 (43) International Publication Date: 9 May 1996 (09.05.96)
(21) International Application Number: PCT/EP95/04117 (22) International Filing Date: 20 October 1995 (20.10.95) (30) Priority Data: 94116558.1 20 October 1994 (20.10.94) EP (34) Countries for which the regional or international application was filed: DE et al. (71) Applicant (for all designated States except US): MORPHOSYS GESELLSCHAFT FÜR PROTEINOPTIMIERUNG MBH [DE/DE]; Frankfurter Ring 193a, D-80807 Munich (DE). (72) Inventors; and (75) Inventors/Applicants (for US only): PACK, Peter [DE/DE]; Franz-Wolter-Strasse 4/III, D-81925 Oberföhring (DE). LU-PAS, Andrei [DE/DE]; Fasanenstrasse 68a, D-82008 Unterhaching (DE). (74) Agent: VOSSIUS & PARTNER; P.O. Box 86 07 67, D-81634 Munich (DE).		(81) Designated States: JP, US, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report.</i> (88) Date of publication of the international search report: 13 February 1997 (13.02.97)
(54) Title: TARGETED HETERO-ASSOCIATION OF RECOMBINANT PROTEINS TO MULTI-FUNCTIONAL COMPLEXES (57) Abstract The present invention relates to a method for targeted assembly of distinct active peptide or protein domains into a single complex and to such complexes. The invention relates particularly to the fusion of peptide or protein domains to complementary association domains which are derived from a single tertiary or quaternary structure by segmentation. The association domains are designed to assemble in a complementary fashion, thereby providing multifunctional (poly)peptides.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 95/04117

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 C12N15/11 C07K14/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C12N C07K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	EP,A,0 404 097 (BEHRINGWERKE) 27 December 1990 see the whole document ---	1
A	WO,A,94 20140 (THE USA GOVERNMENT) 15 September 1994 see the whole document ---	1
A	WO,A,93 15210 (MERCK PATENTS) 5 August 1993 see the whole document ---	1
A	WO,A,93 11162 (PROTEIN DESIGN LABS.) 10 June 1993 see the whole document ---	1
	--- -/--	



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *Δ* document member of the same patent family

Date of the actual completion of the international search

17 April 1996

Date of mailing of the international search report

02.05.96

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+31-70) 340-3016

Authorized officer

Masturzo, P

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 95/04117

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>P PACK: "Dissertation zur Erlangung des Doktorgrades der Fakultät für Chemie und Pharmazie der Ludwig-Maximilians-Universität München: Mini-Antikörper. Bivalente, tetravalente und bispezifische Immunglobuline aus E. coli " 22 April 1994 , MUNICH, GERMANY XP002000449 cited in the application see the whole document -----</p>	1
A	<p>WO,A,92 10209 (WISTAR INSTITUTE) 25 June 1992 see the whole document -----</p>	1

INTERNATIONAL SEARCH REPORT

International application No.

PCT/EP 95/ 04117

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
Remark: Due to the unclear wording of the main claims and the rather theoretical nature of most examples provided, a complete search was deemed impossible for economical reasons. The search has limited to the subject matter of claim 1 as further limited by claims 2 and 4.
2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 95/04117

Patent document cited in search report	Publication date	Patent family member(s)		Publication date
EP-A-404097	27-12-90	DE-A-	3920358	17-01-91
		AU-B-	639241	22-07-93
		AU-B-	5762190	03-01-91
		CA-A-	2019559	22-12-90
		JP-A-	3048699	01-03-91

WO-A-9420140	15-09-94	AU-B-	6399594	26-09-94
		EP-A-	0688228	27-12-95

WO-A-9315210	05-08-93	AU-B-	3410093	01-09-93
		CA-A-	2128511	05-08-93
		CZ-A-	9401757	15-12-94
		EP-A-	0654085	24-05-95
		HU-A-	68798	28-07-95
		JP-T-	7503366	13-04-95
		NO-A-	942750	13-09-94

WO-A-9311162	10-06-93	AU-B-	3147293	28-06-93
		CA-A-	2123894	10-06-93
		EP-A-	0618929	12-10-94
		JP-T-	7501698	23-02-95

WO-A-9210209	25-06-92	CA-A-	2097060	05-06-92
		EP-A-	0563214	06-10-93
